



Rotator Cuff Repair (Large to Massive) Rehabilitation Guideline

This rehabilitation program is designed to return the individual to their activities as quickly and safely as possible. It is designed for rehabilitation following large to massive rotator cuff repairs. Modifications to this guideline may be necessary depending on physician-specific instruction, size and location of tear, tendons involved, acute vs. chronic condition, length of time immobilized, age, first vs. revision, pre-morbid function, tissue quality, fatty infiltration and atrophy, smoking, hypercholesterolemia chronicity of shoulder symptoms, higher levels of cuff retraction and diabetes. This evidence-based large to massive rotator cuff repair physical therapy guideline is criterion-based. Time frames and visits in each phase will vary depending on many factors including patient demographics, goals and individual progress. This guideline is designed to progress the individual through rehabilitation to full sport and activity participation. The therapist may modify the program appropriately depending on the individual's goals for activity.

This guideline is intended to provide the treating clinician with a frame of reference for rehabilitation. It is not intended to substitute clinical judgment regarding the patient's post-operative care based on exam or treatment findings, individual progress and/or the presence of concomitant procedures or post-operative complications. If the clinician should have questions regarding post-operative progression, they should contact the referring physician.

General Guidelines/Precautions:

- Patients with poor compliance to post-operative restrictions in the first 6 weeks show a 152x higher risk of re-tear or non-healing repair. Poor compliance at 12 weeks has 7x risk and poor compliance at week 26 has a 39x risk.
- Immediate post-operative precautions expected
 - No movements beyond neutral extension
 - Keep pillow or towel roll under the arm when lying on back
 - Patient should always be able to see their elbow
 - No reaching behind the back
 - No lifting, pulling or pushing of objects with the involved upper extremity
 - No pushing off with the involved upper extremity during transfers
 - No active range of motion
 - No aggressive, painful passive range of motion or stretching that promotes muscle over-activity or spasm
- Immobilization generally for 6-8 weeks per physician approval
- Protected PROM during the first 6-8 weeks
- Consider Blood Flow Restriction (BFR) training based on patient tolerance. Please see BFR guideline for further information.
- AROM initiated at 8 weeks within the range that shows good mechanics and no pain (weight of arm only)
- Strengthening initiated at week 12
- Return to sport (generally 6-9 months)
 1. Physician approval
 2. Full ROM
 3. Strength within 10% of contralateral side
 4. Shows confidence with sport-specific training with pain 0-2/10.
 5. Independent strength program recommended for at least one year post-surgery
- Anatomic failure is associated with increasing age, poor tissue quality, fatty infiltration, atrophy, smoking, hypercholesterolemia and diabetes
 1. Anatomic failure tends to occur in the first 3-6 months.
- Special considerations that are not accounted for in the guideline below:
 - o Subscapular repair
 1. 0-4 weeks: ER to neutral
 2. 4-6 weeks: Gentle passive ER from neutral to patient tolerance
 3. Extension limited to neutral for 6 weeks
 4. 6+ weeks: Gentle stretching into ER
 5. No resisted IR for 12 weeks
 - o Biceps tenodesis
 6. No active elbow flexion for 6 weeks

Rotator Cuff Repair (Large to Massive) Rehabilitation Guideline (6-9 months to expected D/C)

PHASE	SUGGESTED INTERVENTIONS	GOALS/MILESTONES FOR PROGRESSION
<p>Phase I <i>Patient Education Phase</i></p> <p>Prior to surgery</p>	<p>Discuss: Anatomy, existing pathology, post-op rehab schedule, bracing, precautions and expected progressions</p> <p>Immediate Post-Operative Instructions:</p> <ul style="list-style-type: none"> • Pendulum hang position • Pendulum forward/back and side-to-side with <7-inch arc may be initiated at 2 weeks • Elbow, wrist and hand AROM with no weight 	<p>Goals of Phase:</p> <ol style="list-style-type: none"> 1. Improve ROM and strength to tolerance prior to surgery. 2. Educate patient on appropriate expectation framework for post-operative rehabilitation <p>Criteria to Advance to Next Phase:</p> <ol style="list-style-type: none"> 1. Progress to Phase II post-operatively
<p>Phase II <i>Maximum Protection Phase</i></p> <p>0-4 Weeks</p> <p>Expected visits: 2-4</p>	<p>Discuss: Anatomy, existing pathology, post-op rehab schedule, bracing, precautions, posture and expected progressions</p> <p>Specific Instructions:</p> <ul style="list-style-type: none"> • Formal therapy to start 2-4 weeks post-op • Patient may be instructed on scapular movements and pendulums at a follow-up visit with physician • No movements beyond neutral extension • No reaching behind the back • No lifting, pulling or pushing including during transfers • No AROM of involved shoulder • No aggressive, painful PROM or stretching <p>Suggested Treatments: Modalities as indicated: Edema- and pain-controlling treatments as needed</p> <p>Range of motion:</p> <ul style="list-style-type: none"> • AROM: <ul style="list-style-type: none"> - Neck, elbow, wrist and hand - Scapular retraction/depression to neutral (elbow not behind the back) - Active thoracic extension • PROM <ul style="list-style-type: none"> - Passive pendulum: forward/back, side/side. Less than 7-inch arc - Passive elevation to patient's tolerance up to 90 degrees. - Therapist-assisted passive ER in supine with arm in 20 degrees abduction - Gentle, passive, pain-free supine IR in the plane of the scapula to 30 degrees • Manual therapy <ul style="list-style-type: none"> - Can initiate grade I-II glenohumeral mobs in the plane of the scapula. Directions include posterior, anterior and long-axis traction - Thoracic PA mobilizations can be done seated in weeks 1-2. Can be done prone in weeks 2-4 if tolerated. <p>Other Activities: HEP prescription</p>	<p>Goals of Phase:</p> <ol style="list-style-type: none"> 1. Protect repair 2. Prevent contractures above and below the shoulder joint 3. Manage inflammation and pain 4. Gradual improvements in passive range of motion per guidelines <p>Criteria to Advance to Next Phase:</p> <ol style="list-style-type: none"> 1. Controlled post-operative pain 2. PROM of forward elevation 60-90 degrees in scapular plane 3. PROM of ER in scapular plane: 20 degrees <p style="text-align: right;"><i>(continued on next page)</i></p>

<p>Phase III</p> <p><i>Healing/Protective Phase</i></p> <p>Weeks 4-6</p> <p>Expected visits: 2-4</p>	<p>Specific Instructions:</p> <ul style="list-style-type: none"> • Continue with previous exercise program • Continue sling use unless resting at home • Continue precautions from previous phase • Avoid a head forward, rounded shoulder posture and promote thoracic extension <p>Suggested Treatments:</p> <ul style="list-style-type: none"> • PROM <ul style="list-style-type: none"> - Initiate self-assisted passive ER with a stick upright and/or supine. If supine, limit extension with towel roll behind elbow on table - Therapist-assisted flexion PROM in supine - Progress supine passive ER with stick from 30 degrees to 60 degrees abduction per tolerance • AROM/Strength <ul style="list-style-type: none"> - Scapular retraction and depression AROM - Elbow, wrist and hand AROM - Thoracic extension AROM - Submaximal pain-free elbow flexion and extension isometrics with arm against the body to not resist against shoulder elevation • Manual therapy <ul style="list-style-type: none"> - Grade I and II joint mobs may be used for pain relief/relaxation (GH, AC, ST, SC) - Thoracic PA mobs as needed: Seated or supine to tolerance <p>Exercise Examples:</p> <ul style="list-style-type: none"> • Passive pendulum • Self-assisted passive, pain-free ER with a stick supine or upright • Scapular retraction 	<p>Goals of Phase:</p> <ol style="list-style-type: none"> 1. Protect repair 2. Prevent contractures above and below the shoulder joint 3. Manage inflammation and pain 4. Gradual improvement in PROM per guidelines 5. Toleration of progressed exercise program 6. Passive ER in plane of the scapula: 45 degrees 7. Passive ER at 60 degrees abduction: 45 degrees 8. Passive shoulder flexion 90-120 degrees <p>Criteria to Advance to Next Phase:</p> <ol style="list-style-type: none"> 1. Appropriate healing of the repair by adherence to precautions, immobilization guidelines and exercise protocol 2. Manageable pain level
---	---	--

(continued on next page)

<p>Phase IV <i>Minimal Protection/ Mobility Phase</i></p> <p>Weeks 6-12</p> <p>Expected visits: 6-15</p>	<p>Specific Instructions:</p> <ul style="list-style-type: none"> • Avoid performing activities over shoulder height • Avoid sudden or ballistic movements • No aggressive strengthening • Avoid lifting, pulling or pushing of objects • Weaning from sling according to physician guidelines <p>Suggested Treatments:</p> <ul style="list-style-type: none"> • PROM <ul style="list-style-type: none"> - Initiate self-assisted passive chair roll out flexion - to tolerance (6 weeks) - Continue ER stretching from 30-90 degrees abduction - Initiate shoulder extension and abduction to tolerance (7 weeks) - At 10 weeks, initiate gentle IR stretching including behind the back and in abduction by week 12 • Manual therapy <ul style="list-style-type: none"> - Grade III-IV glenohumeral/scapulothoracic mobilizations for mobility as necessary • AAROM and AROM <ul style="list-style-type: none"> - Active warmup can be done prior to PT via UBE and or active ER/IR in plane of the scapula gravity minimized (8 weeks) - Initiate upright AAROM into flexion and scaption (pulleys or self-assisted) (8 weeks) - As quality of movement improves, progress flexion/scaption from AAROM to AROM, beginning with short lever or gravity modified positions (supine or reclined) - Progress ER AROM from upright to side-lying to tolerance <p>Exercise Examples:</p> <ul style="list-style-type: none"> • Active warmup with non-resisted UBE, pendulum or active upright internal/external rotation • PROM and low load/long duration passive stretching into ER in varying degrees of abduction, into flexion and into scaption. • AAROM/AROM short arc motions in newly acquired range of motion • Shoulder extension ROM to tolerance • Pulleys or wall walks (thumb up with assist of contralateral arm when above 90 degrees) • Gentle IR behind the back (10 weeks) • Progression from AAROM to AROM in flexion as quality of movement improves • As quality of movement improves, initiate and progress AROM endurance training in flexion, scaption, IR and ER <ul style="list-style-type: none"> - Progress from 10 to 30 reps, 1-3 sets 1x/day, 3x/week as tolerated - Endurance work should be in a pain-free arc that avoids substitution patterns - Progress ER from upright to side-lying AROM • Scapular exercise 10 weeks <ul style="list-style-type: none"> - Inferior glide isometric: Shoulder girdle depression while hand rests comfortably on a table - Low row isometric: Scapular depression with extension near neutral • Scapular exercises: 10-12 weeks, 0-light resistance <ul style="list-style-type: none"> - Row, supine protraction, prone extension, scapular clock, side-lying external rotation with scapular setting and no external resistance • Sub-max pain-free GH isometrics at 10 weeks <ul style="list-style-type: none"> - Flexion, Abduction/Scaption, IR, ER in the neutral position • Isotonics <ul style="list-style-type: none"> - Supported biceps and triceps (8 weeks) - Progress to unsupported biceps/triceps at 10 weeks • Rhythmic stabilization progression: <ul style="list-style-type: none"> - Supine ER/IR in the neutral position (8 weeks) - Supine flexion/extension 90 degrees (10-12 weeks) - Ball on table (10-12 weeks) • Cardiovascular training week 8-walking, stationary bike, elliptical without arms 	<p>Goals of Phase:</p> <ol style="list-style-type: none"> 1. Preserve integrity of the repair 2. Able to tolerate initiation and progression of active shoulder flexion and scaption without compensatory hiking 3. Restore functional PROM in all planes with normal movement patterns 4. Decrease pain and inflammation 5. Able to tolerate initiation of submaximal, pain-free muscle activation exercises <p>Criteria to Advance to Next Phase:</p> <ol style="list-style-type: none"> 1. PROM total arc and flexion within 10 degrees of contralateral side 2. AROM shows no substitution patterns, appropriate scapulothoracic rhythm and minimal (NPRS 0-2/10) to no pain in available range of motion
---	--	--

(continued on next page)

<p>Phase V <i>Strengthening and Proprioceptive Phase</i></p> <p>Weeks 12+</p> <p>Expected visits: 5-12</p>	<p>Specific Instructions:</p> <ul style="list-style-type: none"> • Avoid sudden lifting, jerking, pushing or pulling movements • No uncontrolled movements • Avoid heavy lifting, especially above shoulder height (weight lifted must not cause pain or compensatory hiking) <p>Suggested Treatments:</p> <ul style="list-style-type: none"> • Active warmup • Continue with ROM and stretching as needed • Continue biceps and triceps strengthening • Continue proprioception and kinesthetic awareness • Scapulothoracic, glenohumeral, rotator cuff strengthening • Pain management based on education on appropriate progression of activity <p>Exercise Examples:</p> <p>Flexibility:</p> <ul style="list-style-type: none"> • Continue with end-range stretching and manual therapy as needed to restore full total arc and flexion ROM <p>Strength:</p> <ul style="list-style-type: none"> • Biceps curls, triceps press-down • 30-30 ER and IR • Row • Supine serratus punch or dynamic hug • Shoulder flexion, initially only to 90 • Prone or bent over horizontal abduction in external rotation • Scaption, initially to 90 degrees • Straight arm row <p>Exercises that can be added at 18 weeks</p> <ul style="list-style-type: none"> • 90-90 ER and IR in overhead athletes • Prone scaption • Progression to overhead flexion and scaption as tolerated in absence of impingement symptoms or substitution patterns • PNF patterns • Advance CKC exercises over time from partial- to full weight-bearing exercises <p>Proprioception and kinesthetic awareness:</p> <ul style="list-style-type: none"> • Rhythmic stabilization, ball on wall, body blade, etc. 	<p>Goals of Phase:</p> <ol style="list-style-type: none"> 1. Facilitate and maintain functional ROM and quality of movement 2. Tolerate progression of program for muscular strength, power and endurance. <p>Criteria to Advance to Next Phase:</p> <ol style="list-style-type: none"> 1. Strength: 4+/5 to 5/5 or 75%-90% of contralateral side with hand-held dynamometer tested at 22-24 weeks 2. Full ROM in all planes with normal movement mechanics 3. Pain free with basic ADLs and Phase V strengthening 4. Quick DASH <10% disability
<p>Phase VI <i>Advanced Movement and Impact Phase</i></p> <p>Months 6-9 months</p>	<p>Specific Instructions:</p> <ul style="list-style-type: none"> • With overhead athletes, initiate Phase III progressing to Phase IV of Overhead Athlete Rehab Guideline • Initiate Sanford Interval Throwing Program 	<p>Return to Sport:</p> <ul style="list-style-type: none"> • Orthopedic approval • Full, non-painful ROM with no compensatory mechanisms • Strength: MMT 5/5 or 90% of contralateral side with hand-held dynamometer or isokinetic machine • Special considerations for overhead athletes: <ul style="list-style-type: none"> - Successful progression of interval throwing program to 180 feet with no pain. - Consider throwing mechanics assessment - ER/IR Ratio >80% - Hand-held dynamometry at 90-degree abduction <p>Outcome measure options:</p> <ul style="list-style-type: none"> - Quick DASH - Kerlan-Jobe - Patient-Specific Functional Scale - FOTO - See Upper Extremity Testing guideline for further functional testing options depending on activity level of the patient

****NOTE: Progression of functional activities should be performed only as pain and proper biomechanics allow. Emphasis should be on proper mechanics and limiting compensatory mechanisms with exercises and activities.**

REFERENCES:

1. Cools AM, Dewitte V, Lanszweert F, et al. Rehabilitation of scapular muscle balance: Which exercises to prescribe. *The American Journal of Sports Medicine*. 2007; 35: 1744-1751.
2. Davies GJ, Ellenbecker TS. Focused exercise aids shoulder hypomobility. *Biomechanics*: 1999; Nov:77-81.
3. Decker MJ, Hintermeister RA, Faber KJ et al. Serratus anterior muscle activity during selected rehabilitation exercises. *American Journal of Sports Medicine* 1999; 27: 784-791
4. Decker MJ, Tokish JM, Ellis HB et al. Subscapularis muscle activity during selected rehabilitation exercises. *American Journal of Sports Medicine*. 2003; 31: 126-134
5. Dockery ML, Wright TW, LaStayo PC. Electromyography of the shoulder: An analysis of Passive Modes of Exercise. *Orthopedics*. 1998; 21:1181-1184.
6. Gaunt BW, McCluskey GM, Uhl TL. An electromyographic evaluation of subdividing active-assistive shoulder elevation exercises. *Sports Health: A Multidisciplinary Approach*. 2010; 2: 424-432
7. Ghodadra NS, Provencher MT, Verma NN. Open, mini-open, and all-arthroscopic rotator cuff repair surgery: Indications and implications for rehabilitation. *Journal of Orthopedic and Sports Physical Therapy*. 2009; 39: 81-89.
8. Hatakeyama Y, Itoi E, Pradhan RL. Effect of arm elevation and rotation on the strain in the repaired rotator cuff tendon a cadaveric study. *American Journal of Sports Medicine*. 2001; 29: 788-794
9. Kibler WB, Livingston B. Closed chain rehabilitation for upper and lower extremities. *Journal of the American Academy of Orthopaedic Surgeons*. 2001; 9:412-421.
10. Kibler BW, Sciascia AD, Uhl TL et al. Electromyographic analysis of specific exercises for scapular control in early phases of shoulder rehabilitation. *American Journal of Sports Medicine*. 2008; 36: 1789-1798.
11. Koo SS, Burkart SS. Rehabilitation following arthroscopic Rotator Cuff Repair. *Clinical Sports Medicine*. 2010; 29: 203-211.
12. Koo SS, Parsley BK, Burkhart SS, Schoolfield JD. Reduction of postoperative stiffness after arthroscopic rotator cuff repair: Results of a customized physical therapy regimen based on risk factors for stiffness. *Arthroscopy*. 2011; 27: 155-160.
13. Long JL, Ruberte Thiele RA, Skendzel JG et al. Activation of the shoulder musculature during pendulum exercises and light activities. *JOSPT*. 2010; 40: 230-237
14. Ludewig PM, Reynolds JF. The Association of scapular kinematics and glenohumeral joint pathologies. *Journal of Orthopedic and Sports Physical Therapy*. 2009; 39: 90-104.
15. McCann PD, Wootten ME, Kadaba MP, et al. A kinematic and electromyographic study of shoulder rehabilitation exercises. *Clinical Orthopaedics and Related Research*. 1993; 258: 179-188
16. McMullen J, Uhl TL. A kinetic chain approach of shoulder rehabilitation. *Journal of Athletic Training*. 2000; 35: 329-337.
17. Mey KD, Cagnie B, Van De Velde A, et al. Trapezius muscle timing during selected shoulder rehabilitation exercises. *Journal of Orthopedic and Sports Physical Therapy*. 2009; 39: 743-752.
18. Millett PJ, Wilcox RB, O'Holloran JD. Rehabilitation of the rotator cuff: An evaluation- based approach. *Journal of American Academy of Orthopaedic Surgeons*. 2006; 14: 599-609.
19. Mosley JB, Jobe FW, Pink M. EMG analysis of the scapular muscles during a shoulder rehabilitation program. *American Journal of Sports Medicine*. 1992;20: 128-134.
20. Muraki T, Aoki M, Uchiyama E et al. Strain on the repaired supraspinatus tendon during manual traction and translation glide mobilization on the glenohumeral joint: A cadaveric biomechanics study. *Manual Therapy*. 2007; 12: 231-239.
21. Reinold MM, Escamilla R, Wilk KE. Current concepts in the scientific and clinical rationale behind exercises for glenohumeral and scapulothoracic musculature. *Journal of Orthopedic and Sports Physical Therapy*. 2009; 39: 105-117.
22. Reinold MM, Wilk KE, Fleisig GS et al. Electromyographic analysis of the rotator cuff and deltoid musculature during common shoulder external rotation exercises. *Journal of Orthopedic and Sports Physical Therapy*. 2004; 34: 385-394.
23. Smith J, Dahm DL, Kenton RK, et al. Electromyographic activity in the immobilized shoulder girdle musculature during scapulothoracic exercises. *Archives of Physical Medicine & Rehabilitation*. 2006; 87:923-927.
24. Thigpen CA, Shaffer MA, Gaunt BW, et al. The American Society of Shoulder and Elbow Therapists' consensus statement on rehabilitation following arthroscopic rotator cuff repair. *J Shoulder Elbow Surg*. 2016; 25: 521-535.
25. Townsend H, Jobe FW, Pink M. Electromyographic analysis of the glenohumeral muscles during a baseball rehabilitation program. *American Journal of Sports Medicine*. 1991; 19: 264-272
26. Trenerry K, Walton JR, Murrell GAC. Prevention of shoulder stiffness after rotator cuff repair. *Clinical Orthopaedics and Related Research*. 2005; 430: 94-99.
27. Wilk KE, Crockett HC, Andrews JR. Rehabilitation after rotator cuff surgery. *Techniques in Shoulder & Elbow Surgery*. 2000; 1: 128-144.
28. Wilk KE, Meister K, Andrews JR. Current Concepts in rehabilitation of the overhead throwing athlete. *The American Journal of Sports Medicine*. 2002; 30: 136-151.

In designing the current protocol, the following protocols were reviewed:

1. Gunderson Lutheran Sports Medicine (George Davies)
2. Moon Shoulder Group Vanderbilt University
3. Brigham and Women's Hospital Department of Rehabilitation Services
4. The American Society of Shoulder and Elbow Therapists Arthroscopic Rotator Cuff Repair Rehabilitation Guide
5. Fowler Kennedy

Revision Dates: 2/2017, 08/2021, 04/2025



sanfordhealth.org